

# Transport

**GHP INSECTICIDE**

To control listed Household Pests indoors and around the exterior perimeter of residential, institutional, public, commercial, industrial buildings, and food/feed handling establishments.

<b>EPA Reg. No. 8033-96-279</b>	<b>EPA Est. No. 279-NY-1</b>
<b>Active Ingredient:</b>	<b>By Wt.</b>
Acetamiprid .....	22.73 %
Bifenthrin* .....	27.27 %
<b>Other Ingredients:</b> .....	<u>50.00 %</u>
	100.00%

\*Cis isomers 97% minimum, trans isomers 3% maximum.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

ACCEPTED  
FOR REGISTRATION

NOV 06 2013

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

Classified for  
"RESTRICTED USE"  
in New York State  
under 6NYCRR Part 326

Doc ID: 536899



FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia PA 19103

**Net Weight: 0.3 ounce water soluble bags**

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency Assistance.	
<b>NOTE TO PHYSICIAN</b>	
This product contains a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.	
For Information Regarding the Use of this Product Call 1-800-321-1FMC (1362).	

**PRECAUTIONARY STATEMENTS  
Hazards to Humans (and Domestic Animals)  
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

All pesticide handlers (mixers, loaders and applicators) must wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves. After the product is diluted in accordance with label directions for use, and/or when mixing and loading using a closed spray tank transfer system (such as U-Turn®), or an in-line injector system, shirt, pants, socks, shoes and water-proof gloves are sufficient. In addition, all pesticide handlers must wear a respiratory protection device when working in a non-ventilated space. All pesticide handlers must wear protective eyewear when working in non-ventilated space

Use one of the following NIOSH approved respirator with any R, P or HE filter or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE prefilter.

When treating adjacent to an existing structure, the applicator must check the area to be treated, as well as immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean up is completed.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This pesticide is extremely toxic to wildlife, fish and aquatic invertebrates. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

### Physical and Chemical Hazards

Do not apply water-based dilutions of Transport GHP Insecticide to electrical conduits, motor housings, junction boxes, switch boxes or other electrical equipment because of possible shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Keep water-soluble packets in this container and store in a cool dry place but not below freezing (32°F). Do not remove packets from container except for immediate use.

For the following public health pests, do not apply less than the application rates specified on the label: Ants (including Red Imported Fire Ants and Carpenter Ants), Bed Bugs, Bees, Biting Flies, Carpenter Bees, Centipedes, Chiggers, Clover Mites, Cockroaches, Fleas, Flies, Gnats, Ground-nesting (solitary) bees and wasps, Midges, Mosquitoes, Scorpions, Spider Mites, Spiders (including Black Widow and Brown Recluse), Ticks (including Brown Dog Ticks), Wasps.

### STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Keep out of reach of children and animals. Store in original containers only. Do not allow inner bags to become wet during storage. Do not handle inner bag with wet bare hands or wet gloves. Do not put concentrate or diluted material into food or drink containers.

**In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call Chemtrec at 1-800-424-9300.**

**To Confine Spill:** If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal:** Pesticide wastes are toxic. Do not contaminate water, food or feed by storage or disposal. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. Dispose of excess or waste pesticide by use according to label directions, or contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal (for outer packaging and WSP bags/envelopes):** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available.

### APPLICATION RATE CHART

Application Rate: Number of Transport GHP Water-Soluble Packets	Gals of Water	Concentration of Active Ingredient
1 packet (0.3 oz)	1 gallon	0.11%Active Ingredient

### Dilution Chart

#### Dilution Chart for Perimeter Barrier Applications Around Structures

Application Volume per 1,000 sq. ft.	Transport GHP 0.3 ounce water soluble bags to add			
	Total Mix volume			
	1 Gallon	3 Gallons	5 Gallons	10 Gallons
1.0 gal/1,000 sq. ft	1 packet	3 packets	5 packets	10 packets
2.0 gal/1,000 sq. ft	—	—	—	5 packets
2.5 gal/1,000 sq. ft	—	—	2 packets	—
3.0 gal/1,000 sq. ft	—	1 packet	—	—
5.0 gal/1,000 sq. ft	—	—	1 packet	2 packets

### Foam Applications

The Transport GHP Insecticide dilution may be converted to foam with expansion characteristics from 2 to 40 times for localized control or prevention of pests including ants, bees, wasps or other arthropods harboring in walls, under slabs or in other void areas.

Depending on the circumstances, foam applications may be used alone or in combination with liquid dilution applications. Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block voids or structural voids, under slabs, stoops, porches, or to the soil in crawlspaces, and other similar voids.

Foam and liquid application must be consistent with volume and active ingredient instructions in order to insure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the labeled liquid dilution volume of product must be applied, with the remaining percent delivered to appropriate areas using foam application. Refer to label and use recommendations of the foam manufacturer and the foaming equipment manufacturer.

Foam applications are generally a good supplement to liquid treatments in difficult areas, but may be used alone in difficult spots.

Use dry foam (20:1 or greater expansion ratio) for applications to wall voids and stud walls.

Use wet foam (5:1 or lower expansion ratio) for applications to soil, including applications to filled porches.

#### Mixing table for Transport GHP Insecticide Foam

Desired Foam Expansion Ratio	Amount of Transport GHP	Gallons of Water	Finished Foam (Gallons)	Finished Foam % a.i.
4:1	4 packets	1.0	4	0.11
10:1	10 packets	1.0	10	0.11
20:1	20 packets	1.0	20	0.11

### Household Pest Control – Outdoor Pinstream, Spot, Crack and Crevice or Perimeter Treatment

Controls	Ants (including Red Imported Fire Ants and Carpenter Ants) (but excluding Pharaoh ants and Argentine ants)	Ground-nesting (solitary) bees and wasps
	Bees	Midges
	Beetles*	Millipedes
	Biting Flies	Mosquitoes
	Boxelder Bugs	Moths
	Centipedes	Pillbugs
	Chiggers	Scorpions
	Clover Mites	Silverfish
	Cockroaches	Sowbugs
	Crickets	Spider Mites
	Earwigs	Spiders (including Black Widow)
	Elm Leaf Beetles	Springtails
	Firebrats	Stink Bugs
	Fleas	Ticks (including Brown Dog Ticks)
	Flies	Wasps
	Gnats	*Not for use in California

**Where to Apply**  
Apply as a pinstream, spot, crack and crevice, or perimeter spray on and around outside surfaces of residential and non-residential buildings and structures including, exterior siding, foundations, porches, window frames, eaves, patios, garages, areas adjacent or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, storage sheds, and other structures, and other areas where pests congregate or have been seen. While this product is not for use on turf or lawns, adjacent (i.e., perimeter) treatment is permitted, as directed by the Perimeter Treatment Application Rate. Follow Additional Application Restrictions for Residential Outdoor Surface and Space Sprays.

**Perimeter Treatment Application Rate**  
As a perimeter treatment, apply as a continual band up to 10 foot wide around the structure and upwards along the foundation to a height of up to 3 feet and around windows, doors, other penetrations and roof eaves, soffits, and overhangs.

Apply 0.11% concentration of Transport GHP Insecticide in sufficient amount of water (see Dilution Chart) and apply adequately to cover 1,000 square feet. Dilutions may be applied at either high or low volumes. Do not apply more than 0.11% finished dilution per 1,000 square feet.

**Mixing Directions**  
Fill tank 1/4 to 1/3 full with water. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose. Add Transport GHP Insecticide bag(s). Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes.

For backpacks and handheld sprayers, use the 0.3 ounce sized bag. Fill the tank 1/2 full with water. Add Transport GHP Insecticide water soluble bag(s). Agitate tank after adding water and allow 1 to 2 minutes for bags to dissolve in water. Add remaining water. Close application equipment and agitate or shake tank to dissolve any bag remnants in the tank.

<b>Mixing Directions (Continued)</b>	Transport GHP Insecticide may also be mixed into full tanks of water, but requires substantial agitation to insure uniformity of the dilution.
<b>Repeat Applications</b>	Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity.
<b>Restrictions</b>	<p>Do not water the treated area to the point of run-off. Do not make applications during rain.</p> <p>All outdoor applications, if permitted elsewhere on this label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label: 1) Applications to soil or vegetation, as listed on this label, around structures; 2) Applications to the side of a building, up to a maximum height of 3 feet above grade; 3) Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure; 4) Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width; 5) Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests.</p> <p>Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, drive-ways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.</p> <p>Applications to vertical exterior surfaces (e.g., foundations are permitted to a maximum height of 3 feet from ground level. Sections of vertical exterior surfaces that abut non-porous horizontal surfaces can only be treated if either 1) these sections are protected from rainfall and spray from sprinklers or 2) they do not drain into a sewer, storm drain, or curbside gutter (e.g., not to sections that abut driveways or sidewalks that drain into streets).</p> <p>For sections of foundation that abut non-porous horizontal surfaces, the treated areas must be protected from rainfall and spray from sprinklers so they do not drain into a sewer, storm drain, or curbside gutter (e.g. not to sections that abut driveways or sidewalks that drain into streets).</p> <p>Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.</p>

<b>Carpenter Ants in trees, utility poles, fencing, deck materials and similar structural members</b>	Drill to locate the interior infested cavity and inject or foam the 0.11% dilution (see Dilution Chart) into the cavity using a sufficient volume and an appropriate treatment tool with a splash-back guard.
<b>Ground-nesting (solitary) Bees and Wasps (including Bumble Bees, Sweat Bees, Mining Bees, Digger Bees, Leafcutting Bees, Threadwaisted Wasps)</b>	<p>Ground-nesting bees and wasps are helpful biocontrol agents and valuable pollinators. They should be ignored if possible. If control is necessary, however, nest entrances must first be located. Watch the insects during the day when they are active. Groups of single nests occur in bare soil, grassy/weedy areas, or cavities of shrubs, stems, twigs, or logs. Treatment of tunnels and the surrounding area at dusk or after dark improves product contact to individual in-ground nest dwellers.</p> <p>Individual nest drenches should be applied by using 0.3 ounces of Transport GHP Insecticide per 1,000 square feet in and around each cavity. Cover the entrance hole with soil after application. For preventative treatment, broadcast Transport GHP Insecticide in enough finished volume of water to penetrate the groundcover so that 0.3 ounces are applied per 1,000 Square Feet.</p>
<b>Wood piles and stored lumber</b>	To protect firewood or lumber from carpenter ants, make up a 0.11% dilution (see Dilution Chart) of Transport GHP Insecticide and apply as a spot treatment to the soil beneath where the firewood or lumber will be stacked at the rate of one gallon of dilution per 8 square feet. Treated wood can be burned as firewood or used as lumber one month after treatment.
<b>Underground Services</b>	<p>Underground Services such as: wires, cables, utility lines, pipes, conduits, etc. Services may be within structures or located outside of structures.</p> <p>Soil treatment may be made using Transport GHP Insecticide dilution to prevent attack by Termites and Ants.</p> <p>Apply 2 gallons of 0.11% dilution (see Dilution Chart) per 10 linear feet to the bottom of the trench and allow liquid to soak into the soil. Lay services on the treated soil and cover with approximately 2 inches of fill soil. Apply another 2 gallons per 10 linear feet over the soil surface to complete the treatment barrier. In wide trenches, only treat the soil in the area near the services. It is important to establish a continuous barrier of treated soil surrounding the services.</p> <p>Where soil will not accept the above-labeled volume, 1 gallon of 0.11% dilution of Transport GHP Insecticide may be used per 10 linear feet of trench both to the bottom of the trench and over the soil on top of the services.</p> <p>Finish filling the trench with treated fill soil. The soil where each service protrudes from the ground may be treated by trenching/rodding of no more than 1 to 2 gallons of 0.11% dilution into the soil.</p>
<b>Posts, Poles and Other Constructions</b>	<p>Create an insecticidal barrier in the soil around wooden constructions such as signs, fences and landscape ornamentation by applying a dilution.</p> <p>Previously installed poles and posts may be treated by sub-surface injection or treated by gravity-flow through holes made from the bottom of a trench around the pole or post. Treat on all sides to create a continuous insecticidal barrier around the pole. Use 1 gallon of 0.11% dilution (see Dilution Chart) per foot of depth for poles and posts less than six inches in diameter. For larger poles, use 1.5 gallons of 0.11% dilution per foot of depth. Apply to a depth of 6 inches below the bottom of the wood. For larger constructions, use 4 gallons per 10 linear feet per foot of depth.</p>
<b>Pests Under Slabs</b>	Infestations of Arthropods, such as Ants, Cockroaches and Scorpions inhabiting under slab areas may be controlled by drilling and injecting or horizontal rodding and then injecting 1 gallon of 0.11% dilution (see Dilution Chart) per 10 square feet or 2 gallons of 0.11% dilution per 10 linear feet.
<b>Pest Control in Crawlspace and Voids</b>	Apply Transport GHP Insecticide 0.11% dilution (see Dilution Chart) to all surfaces in crawlspace and/or voids to control ants, fleas, roaches, scorpions, or other arthropods. Product may also be applied through insecticidal delivery systems such as piping or flexible tubing mounted under and/or around the structure as a crack and crevice or spot treatment. Treat surfaces to point of runoff. Keep children and pets off surface until dry.

<b>Outdoor Ant Control</b>	
<b>Carpenter Ants</b>	<p>Apply Transport GHP Insecticide as a pinstream, spot, crack and crevice, or perimeter spray to carpenter ant trails around doors and windows and other places where carpenter ants have been observed or are expected to forage. For best results, locate and treat carpenter ant nests. Apply a perimeter treatment using either low or high volume applications described in the Household Pest Control - Outdoor section of this label. The higher dilutions and/or application volumes, as well as more frequent applications, may be necessary when treating concrete surfaces for ant control. The following procedures must be followed to help achieve maximum control of the pest:</p> <ol style="list-style-type: none"> <li>1) Treat non-porous surfaces only in areas protected from rainfall and spray from sprinklers with low volume applications using 0.11% dilution per 1,000 square feet.</li> <li>2) Treat porous surfaces and vegetation with high volume applications using 0.11% dilution per 1,000 square feet.</li> <li>3) Treat the trunks of trees that have carpenter ant trails or upon which carpenter ants are foraging using 0.11% dilution and applying this dilution to thoroughly wet the bark from the base of the tree to as high as possible on the trunk.</li> </ol>
<b>Nuisance Ants Outdoor (including fire ants)</b>	<p>For best results, locate and treat ant nests. Apply Transport GHP Insecticide as a pinstream, spot, crack and crevice or perimeter treatment to ant trails around doors and windows and other places where ants have been observed or are expected to forage. Apply a perimeter treatment described in the Household Pest Control - Outdoor section of this label. More frequent applications, may be necessary when treating concrete surfaces for ant control. The following procedures must be followed to help achieve maximum control of the pest:</p> <ol style="list-style-type: none"> <li>1) Treat non-porous surfaces only in areas protected from rainfall and spray from sprinklers with low volume applications using 0.11% dilution per 1,000 square feet.</li> <li>2) Treat porous surfaces and vegetation with high volume applications using 0.11% dilution per 1,000 square feet.</li> </ol>
<b>Specific Pest Control Applications</b>	
<b>Ant and Fire Ant Mounds</b>	Drench mounds with 1-2 gallons of Transport GHP Insecticide at a 0.11% dilution (see Dilution Chart) to each mound area by sprinkling the mound until it is wet and treat 3-feet around the mound. Use the higher volume for mounds larger than 12". For best results, apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day.

<b>Household Pest Control - Indoor</b>		
<b>Controls</b>	<p>Ants (including Red Imported Fire Ants and Carpenter Ants) (but excluding Pharaoh ants and Argentine ants)</p> <p>Bed Bugs</p> <p>Bees</p> <p>Beetles (including Carpet Beetles)</p> <p>Boxelder Bugs</p> <p>Centipedes</p> <p>Cockroaches</p> <p>Crickets</p> <p>Earwigs</p> <p>Firebrats</p> <p>Fleas</p> <p>Flies</p>	<p>Gnats</p> <p>Midges</p> <p>Millipedes</p> <p>Moths (including Cloth Moths)</p> <p>Pillbugs</p> <p>Scorpions</p> <p>Silverfish</p> <p>Sowbugs</p> <p>Spiders (including Black Widow and Brown Recluse)</p> <p>Springtails</p> <p>Stink bugs</p> <p>Ticks</p> <p>Wasps</p>

<b>Where to Apply</b>	Apply for residual pest control in residential and non-residential buildings and structures. Apply either as a crack and crevice, pinstream, spot, coarse, low-pressure spray (25 p.s.i. or less) or with a paintbrush. Apply to areas where pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, dishwashers, cabinets, sinks, furnaces, stoves, the underside of shelves, drawers and similar areas and other possible pest harborage sites. Do not use as a space spray. Pay particular attention to cracks and crevices. Do not apply as a broadcast spray indoors.
<b>Application Rate</b>	Apply 0.11% concentration of active ingredient in sufficient amount of water (see Dilution Chart) and apply adequately to cover 1,000 square feet. Do not apply more than 0.11% finished dilution per 1,000 square feet.
<b>Mixing Directions</b>	For backpacks and handheld sprayers, use the 0.3 ounce sized bag. Fill the tank 1/2 full with water. Add Transport GHP Insecticide. Allow 1 to 2 minutes for bags to dissolve in water. Add remaining water. Close application equipment and agitate or shake tank to dissolve any bag remnants in the tank. Mixing with warm water (68°F or higher) will decrease water soluble bag dissolution time.
<b>Ants (including Carpenter Ants and Nuisance Ants) (but excluding Pharaoh ants and Argentine ants)</b>	Apply to areas where ants have been observed or are expected to forage. Particular attention should be given to treating entry points into the home or premises such as around doors and windows and other places where ants and ant trails may be found. For added Carpenter Ant control, spray or foam into cracks and crevices or drill holes and spray, mist or foam into voids where Carpenter Ants or their nests are present. When using Transport GHP Insecticide in combination with baits, apply Transport GHP Insecticide as instructed above, and use baits in other areas that have not been treated with Transport GHP Insecticide.
<b>Bed Bugs</b>	To control Bed Bugs, use a 0.11% active ingredient dilution rate where evidence of bed bugs occurs. (See Dilution chart for mixing instructions) Thorough application should be made to crack and crevices where evidence of bed bugs occurs. This includes bed frames, box springs, mattresses, inside empty luggage, dressers and clothes closets and carpet edges, high and low wall moldings and wallpaper edges, wall hangings, mirrors, pictures, electrical switch plates, furniture, door frames, bookcases and windowframes. For infested mattresses, remove linens and wash before reuse. Apply to tufts, seams, folds, and edges until moist. Allow to dry before remaking bed. When treating furniture, pay special attention to tufts, folds, seams and difficult-to-access areas. On furniture do not apply to seating areas, arms or areas where direct skin contact can occur. Do not use this product on bed linens, pillows, or clothes. Remove all clothes and other articles from luggage, dressers or clothes closets before application. Allow all treated areas to thoroughly dry before use. Not recommended for use as sole protection against bedbugs.
<b>Bees and Wasps</b>	To control Bees, Wasps, Hornets, and Yellow Jackets indoors, application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible.
<b>Occasional Invaders</b>	To control Boxelder Bugs, Centipedes, Earwigs, Beetles, Millipedes, Pillbugs, and Sowbugs, apply around doors and windows and other places where these pests may be found or where they may enter premises. Spray baseboards, storage areas and other locations.
<b>Crawling and Flying Insect Pests</b>	To control Cockroaches, Crickets, Firebrats, Flies, Gnats, Midge, Moths, Scorpions, Silverfish, Spiders, apply as a coarse, low pressure spray to areas where these pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under dishwashers, refrigerators, cabinets, sinks, furnaces, and stoves, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices.
<b>Restrictions</b>	In the home, cover all food processing surfaces and utensils during treatment or thoroughly washed before use. Remove or cover exposed food before application. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

## Specific Indoor Pest Control Applications

<b>Warehouses and Stores</b>	Transport GHP Insecticide may be applied as a spot or crack and crevice treatment in non-food storage warehouses and stores. Apply no more than 0.11% dilution (see Dilution Chart) of Transport GHP Insecticide per 1,000 square feet to provide adequate coverage. Apply to all areas that may harbor pests, including under and between pallets, bins and shelves. Do not apply direct to food grain bins (interior) or animals.
------------------------------	---

## Food/Feed Handling Establishment Applications

<b>Controls</b>	<table border="0"> <tr> <td>Ants (including Red Imported Fire Ants and Carpenter Ants) (excluding Pharaoh ants and Argentine ants)</td> <td>Flies</td> </tr> <tr> <td>Bed Bugs</td> <td>Gnats</td> </tr> <tr> <td>Bees</td> <td>Midges</td> </tr> <tr> <td>Beetles (including Carpet Beetles)</td> <td>Millipedes</td> </tr> <tr> <td>Boxelder Bugs</td> <td>Moths (including Cloth Moths)</td> </tr> <tr> <td>Centipedes</td> <td>Pillbugs</td> </tr> <tr> <td>Cockroaches</td> <td>Scorpions</td> </tr> <tr> <td>Crickets</td> <td>Silverfish</td> </tr> <tr> <td>Earwigs</td> <td>Sowbugs</td> </tr> <tr> <td>Firebrats</td> <td>Spiders (including Black Widow and Brown Recluse)</td> </tr> <tr> <td>Fleas</td> <td>Springtails</td> </tr> <tr> <td></td> <td>Stink bugs</td> </tr> <tr> <td></td> <td>Ticks</td> </tr> </table>	Ants (including Red Imported Fire Ants and Carpenter Ants) (excluding Pharaoh ants and Argentine ants)	Flies	Bed Bugs	Gnats	Bees	Midges	Beetles (including Carpet Beetles)	Millipedes	Boxelder Bugs	Moths (including Cloth Moths)	Centipedes	Pillbugs	Cockroaches	Scorpions	Crickets	Silverfish	Earwigs	Sowbugs	Firebrats	Spiders (including Black Widow and Brown Recluse)	Fleas	Springtails		Stink bugs		Ticks
Ants (including Red Imported Fire Ants and Carpenter Ants) (excluding Pharaoh ants and Argentine ants)	Flies																										
Bed Bugs	Gnats																										
Bees	Midges																										
Beetles (including Carpet Beetles)	Millipedes																										
Boxelder Bugs	Moths (including Cloth Moths)																										
Centipedes	Pillbugs																										
Cockroaches	Scorpions																										
Crickets	Silverfish																										
Earwigs	Sowbugs																										
Firebrats	Spiders (including Black Widow and Brown Recluse)																										
Fleas	Springtails																										
	Stink bugs																										
	Ticks																										
<b>Where to Apply</b>	Applications of this product are permitted in both food/feed and non-food areas of food/feed establishments as a spot or crack and crevice treatment. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served, which includes areas for receiving, storing, packaging (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, edible oils, syrups) or food. Service areas where food is exposed and the facility is in operation are also considered food areas. Permitted non-food areas of use include, garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets and storage (after canning or bottling). Permitted use sites include: aircraft (do not use in aircraft cabins), apartment buildings, bakeries, bottling facilities, breweries, buses, cafeterias, candy plants, canneries, dairy product processing plants, food manufacturing plants, food processing plants, food service establishments, granaries, grain mills, grocery stores (do not apply directly to the interior of food grain bins) hospitals, hotels, industrial buildings, laboratories, meat/poultry/egg processing plants, mobile/motor homes, nursing homes, offices, pet stores (do not apply directly to the interior of food grain bins or animals), railcars, restaurants, schools, ships, trailers, trucks, vessels, warehouses and wineries.																										
<b>Application Rate</b>	Apply 0.11% per 1,000 square feet in sufficient amount of water (see Dilution Chart) and apply adequately to cover area. Do not apply more than 0.11% of Transport GHP Insecticide per 1,000 square feet.																										
<b>Spot, Crack and Crevice Application</b>	Spot or crack and crevice applications may be made while the facility is in operation; however, cover or remove food from area being treated. Do not apply directly to food.																										
<b>Mixing Directions</b>	Fill tank 1/4 to 1/3 full with water. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose. Add Transport GHP Insecticide. Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes. For smaller tanks, fill tank 1/2 full with water. Add desired Transport GHP Insecticide water soluble bag(s). Allow 1 to 2 minutes for dissolution of water soluble bags. Close application equipment and agitate or shake tank. Transport GHP Insecticide may also be mixed into full tanks of water, but requires substantial agitation to insure uniformity of the dilution.																										
<b>Foam Applications</b>	Transport GHP Insecticide may be converted to a foam and used to treat void spaces or as a spot spray on vertical horizontal surfaces where visual marking of application is desired. Use of a foaming agent increases a.i. surface contact time on challenging surfaces and provides visual marking of the application. Ensure that the foaming agent is approved for food surface/area contact use. See Foam Application section for mixing table and rates.																										

## RESTRICTIONS

Do not apply in a perimeter treatment to areas beyond 10 feet from the foundation of the structure, unless using as a spot treatment.

Do not use as a space spray.

Do not use in and around the exterior perimeter of commercial barns, stables, paddocks. Do not use in grazing areas, feed lots or other similar areas used for housing, boarding, and/or rearing animals. This product may be used around barns and stables on residential property.

Do not apply by air.

Do not apply as a broadcast spray indoors or as a broadcast spray on lawns and turf.

Do not apply in greenhouses or nurseries.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not apply this product through any kind of irrigation system.

Not for use on sod farm turf, golf course turf, or grass grown for seed.

Do not apply to pets, crops, or sources of electricity.

Do not treat electrically active underground services.

Do not treat areas when food is exposed. Cover or remove food from area being treated. Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.

Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product.

Let surfaces dry before allowing people and pets to contact surfaces.

Prior to applying Transport GHP Insecticide to porous surfaces, test a small area and allow it to dry to be sure no deposits will form.

Application equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol® applicators, may also be used to make crack and crevice, deep harborage, spot and general surface treatments of Transport GHP Insecticide.

Wear protective clothing; unvented goggles, gloves and a respirator approved by NIOSH, when applying to overhead areas or in poorly ventilated or confined areas.

## Conditions of Sale of Warranty and Liability:

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC Corporation or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC Corporation and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC Corporation MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT TO THE EXTENT ALLOWED BY STATE LAW. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC Corporation and buyer assumes the risk of any such use.**

To the extent consistent with applicable law, FMC Corporation or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC Corporation AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC Corporation OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC - Trademark of FMC Corporation

Transport - Trademark of Nippon Soda Co., Ltd